

**MANGANESE COMPOSITIONS FOR
REDUCING/PREVENTING SKIN WRINKLES
AND FINE LINES**

CROSS-REFERENCE TO PRIORITY APPLICATION

[0001] This application claims priority under 35 U.S.C. §119 of FR-00/06373, filed May 18, 2000, hereby expressly incorporated by reference.

CROSS-REFERENCE TO COMPANION APPLICATION

[0002] Our copending application Serial No. _____ [Attorney Docket No. 016800-443], filed concurrently herewith and assigned to the assignee hereof.

BACKGROUND OF THE INVENTION

Technical Field of the Invention:

[0003] The present invention relates to the administration, to individuals in need of such treatment, of cosmetic/dermatological compositions comprising effective amounts of manganese or salts thereof for relaxing and/or slackening cutaneous and/or subcutaneous tissue, notably for reducing/preventing wrinkles and fine lines in the skin.

Description of the Prior Art:

[0004] There is a current trend for women, and even men, to wish to appear young for as long as possible and, consequently, to fade out the signs and marks of aging of the skin, which are especially reflected in wrinkles and fine lines. In this respect, advertizing and the fashion world report products intended to maintain a radiant and wrinkle-free skin for as long as possible, which are signs

compartment (depolarization) plays a part in reducing the potential difference between the outside and the inside and thus renders the cell more excitable.

[0010] Specifically, the depolarization of the transverse tubules (invagination of the cell membrane) which spreads to the longitudinal tubules (sarcoplasmic reticulum) induces the instantaneous release of intracellular calcium by these longitudinal tubules. In the presence of calcium, the contractile proteins of striated muscle show ATPase activity which provides the energy required for the contraction.

[0011] Conversely, relaxation of the striated muscle appears following the binding of ATP to the contractile proteins. The intracellular calcium then re-enters the intracellular compartment and its concentration returns to a value close to 10^{-8} M.

[0012] Moreover, it has been shown that botulinum toxin which was originally used to treat spasms, can act on muscular spasticity states (see A. Blitzer et al., Arch. Otolaryngol. Head Neck Surg., **119**, pages 1018 to 1022 (1993)) and on wrinkles of the glabella, which are the wrinkles between the eyebrows (see J.D. Carrutgers et al., J. Dermatol. Surg. Oncol., **18**, pages 17 to 21 (1992)). Consequently, it is possible to act on the muscular contractile component of wrinkles (in particular on the motor plate which corresponds to the nerve/muscle junction).

[0013] In the peripheral nervous system, the junction between a nerve and a muscle constitutes the neuromuscular plate, upstream of which is the efferent nerve pathway known as the motor neurone. Moreover, the cell membranes of each nerve fiber also comprise many ion channels, and in particular calcium channels, which are capable of allowing the corresponding element to permeate in ionic form, which in this particular case is calcium.

[0014] The important role of calcium and of regulating its intracellular concentration in the phenomena of muscle contraction/relaxation, whether these

THE UNIVERSITY OF CHICAGO

[0015] Regulation of the intracellular calcium concentration is possible only because the efflux of calcium corrects the influx. This can be assured only by an expulsion of the cellular calcium by one or more mechanisms capable of overcoming the electrochemical potential gradient mentioned above.

[0017] In order to better describe the calcium-permeability changes, it is currently common to consider that this permeability corresponds to the opening of membrane-bound calcium channels, these channels being operated by variations in the membrane potential (VOC) or by activation of the membrane-bound receptors (ROC). To date, six VOC types of calcium channel (L, N, T, P, Q and R) have been identified.

SUMMARY OF THE INVENTION

[0019] After numerous clinical tests, it has now unexpectedly and surprisingly been determined that contractile muscle fiber, which is under the direct control of the neuromotor influx, serves an essential function in the formation of wrinkles and that the modulation of the neuromotor influx and the

DETAILED DESCRIPTION OF BEST MODE AND SPECIFIC/PREFERRED EMBODIMENTS OF THE INVENTION

[0022] The simplest experimental technique for demonstrating that a pharmacological agent is capable of inhibiting calcium influx entails preincubating a smooth muscle in a calcium-free physiological solution, depolarizing it in a KCl-rich solution and gradually increasing the calcium concentration in the infusion solution. This elicits an increase in the tension of the muscle, the value of which increases to a maximum as a function of the calcium concentration. When this protocol is repeated in the presence of a substance considered to inhibit calcium influx, as was carried out for the first time with cinnarizine, the contractile responses are inhibited in a dose-dependent manner. A similar concept was applied to describe the action of verapamil on the heart. Verapamil was first

considered a β -blocker, but its action is more complex since it exerts inhibitory action on excitation/contraction coupling. On the papillary muscle, verapamil suppresses the contraction by very slightly modifying the action potential. It is this observation which resulted in verapamil being considered as a calcium antagonist.

[0023] Manganese is a metal which is very widespread at the surface of the earth's crust. It belongs to Group VIIa of the Periodic Table, its atomic number is 25 and its atomic weight is 54.93. Manganese has several valences (1 to 7), the divalent and trivalent forms being those that are the most biologically active.

[0024] Manganese is widely used in the metallurgy industry, in the manufacture of dry batteries and as a colorant.

[0025] Plants are all rich in Mn, particularly seeds (about $7 \mu\text{g/g}$), nuts (about $17 \mu\text{g/g}$) and tea. Fruits (about $1 \mu\text{g/g}$) and vegetables (about $2.5 \mu\text{g/g}$) are less rich, but their level is still very high compared with foods of animal origin (meats: about $0.20 \mu\text{g/g}$, fish: about $0.05 \mu\text{g/g}$).

[0026] Conversely, this metal only exists in trace amounts in animals, particularly humans.

[0027] However, its biological role is very important and, even though the harmful effects of a Mn deficiency have not been determined irrefutably in man, the consequences of deficiencies examined in animals indicate that this metal is involved in many metabolic schemes. However, even today, the knowledge regarding the intimate biochemical mechanisms of Mn remains very fragmented.

[0028] Manganese has been implicated in many metabolic pathways:

- (a) clotting;
- (b) thermogenesis (by its action on the thyroid system);
- (c) immunity, in which manganese appears to be necessary for a proper synthesis of antibodies;
- (d) reproduction, its deficiency, which promotes a reduction in the fertility

[0029] Two properties permit explaining certain of the physiopathological roles of manganese:

[0030] Manganese is a metal which activates numerous enzymes and lectins. It intervenes either as a dissociable element or by forming an integral part of the structure of the enzyme (metalloenzymes).

[0031] The internal signal to activate a cell is often triggered by a modification of the intracytoplasmic calcium concentrations. The calcium binds to proteins (calmoduline) which in turn activate kinases. Calcium serves in the transmission of the nerve influx, in the stimulation of certain secretory cells, it triggers the changes in the shape of platelets at the beginning of their activation, etc.

[0033] In order for a substance to be recognized as a calcium-channel inhibitor, also referred to herein as a calcium antagonist, it must be able to reduce the intracellular calcium concentration or reduce the binding of calcium to intracellular proteins such as, for example, calmoduline, as is especially described, for example, by Galizzi, J.P. et al., Biol. Chem., 262, p. 6947 (1987) or Y. Okamiya et al., Eur. J. Pharmacol., 205, p. 49 (1991) or J.A. Wagner et al., J. Neurosci., 8, p. 3354 (1988) or H.R. Lee et al., Life Sci., 35, p. 721 (1984) or Schoemaker H. and Lauger S., Eur. J. Pharmacol., 111, p. 273 (1985) or I.J. Reynolds et al., J. Pharmacol. Exp. Ther., 237, p. 731 (1986).

[0034] For the purposes of the present invention, a substance is recognized as a relaxer when it elicits a relaxation effect on a contracted muscle tissue and/or exerts an inhibitory effect in an experiment model of nerve/muscle junction (motor plate), in particular, in the model described by W. Steinbrecher in "Electrodes for stimulation and bioelectric potential recording," Ed. Biomerstechnich, pages 96-98 (1988).

[0035] Manganese and the salts thereof fully satisfy these criteria.

[0036] As indicated above, the present invention features influencing calcium channels to relax or slacken vascular tissues, and thus combating wrinkles and fine lines, via administering, to an individual subject in need of such treatment, manganese, whether in ionic form, in the form of a salt or in the form of manganese-rich natural, plant or microorganism, particularly bacterial, extracts.

[0037] Thus, this invention features the administration of compositions comprising an effective amount of manganese and/or at least one of the salts thereof, to relax and/or slacken cutaneous and/or subcutaneous tissue, such compositions containing a physiologically acceptable medium (vehicle, diluent or carrier).

[0038] This invention also features, if necessary formulated into a physiologically acceptable medium therefor, administration of an effective amount of natural, plant or microorganism, particularly bacterial, extracts that are rich in manganese or in manganese salt, to relax and/or slacken cutaneous and/or subcutaneous tissue.

[0039] By the term "manganese salts" are intended organic or inorganic manganese salts.

[0040] Exemplary organic salts according to the invention include manganese carbonate, manganese acetate, manganese citrate, manganese oleate, manganese oxalate, etc.

[0041] And exemplary inorganic manganese salts include the mineral salts, for instance manganese chloride, manganese borate, manganese nitrate, manganese

phosphate, manganese sulfate, etc.

[0042] Moreover, except where otherwise indicated, the term “manganese” means manganese which is not only in ionic form but also in the form of salts or in the form of manganese-rich natural, plant or microorganism, particularly bacterial, extracts.

[0043] By the expression “physiologically acceptable medium” is intended a medium which is compatible with the skin, the scalp and/or mucous membranes.

[0044] The present invention also features formulating an effective amount of manganese values into physiologically acceptable media to provide compositions suited for smoothing the skin, and also for attenuating and/or eliminating the microrelief of the skin.

[0045] The subject compositions are well suited for curatively and/or preventively combating wrinkles and fine lines in the skin.

[0046] The subject compositions are particularly well suited for reducing wrinkles and fine lines in the skin.

[0047] More particularly, the relaxation and/or slackening of the cutaneous and/or subcutaneous tissue corresponds to a muscular relaxation or slackening.

[0048] Thus, the effective amount of manganese which may be administered according to the invention depends on the desired effect and may vary over a wide range.

[0049] To provide an order of magnitude, it is intended, according to the invention, to administer manganese in an amount of from 0.0001 % to 10 % of the total weight of the composition and preferably in an amount of from 0.001 % to 5 % of the total weight of the composition.

[0050] When, according to the invention, a manganese-rich natural, plant or microorganism, particularly bacterial, extract is administered, one skilled in this art can easily adapt the amount of extract such that, in the final analysis, the manganese is administered in the amounts indicated above.

[0051] Exemplary manganese-rich natural extracts according to the

invention include the extracts of nut or extracts of tea.

[0052] The compositions of the invention are intended for cosmetic or dermatological applications. The compositions of the invention are preferably suited for cosmetic applications.

[0053] The regime/regimen according to the invention is cosmetic, since intended to modify a person's appearance.

[0054] The compositions according to the invention may be in any presentation form that is conventional for topical, injectable or oral administration.

[0055] The compositions according to the invention may be administered either locally, i.e., topically, or by subcutaneous and/or intradermal injection or by oral administration.

[0056] Preferably, the subject compositions are topically applied.

[0057] The amounts of the various constituents of the compositions according to the invention are those conventionally included in the fields under consideration and which are appropriate for the various formulations.

[0058] For topical application, the compositions of the invention comprise a medium that is compatible with the skin. These compositions may especially be formulated as aqueous, alcoholic or aqueous/alcoholic solutions, gels, water-in-oil or oil-in-water emulsions having the appearance of a cream or a gel, microemulsions or aerosols, or alternatively in the form of vesicular dispersions containing ionic and/or nonionic lipids. Such compositions are formulated according to the usual methods in the fields under consideration.

[0059] These topical-application compositions may, in particular, constitute a protective or care formulation for the face, the neck, the hands or the body (for example day creams, night creams, sunscreen creams or oils or body milks), a makeup composition (for example a foundation) or an artificial tanning composition.

[0060] When the composition of the invention is an emulsion, the proportion of fatty substance it contains may range from 5% to 80% by weight

and preferably from 5% to 50% by weight relative to the total weight of the composition. The fatty substances and emulsifiers contained in the emulsion are selected from among those conventionally employed in the cosmetic or pharmaceutical field.

[0061] Exemplary fatty substances according to the invention include the mineral oils (petroleum jelly), plant oils (liquid fraction of karite butter) and hydrogenated derivatives thereof, animal oils, synthetic oils (perhydrosqualene), silicone oils (polydimethylsiloxane) and fluoro oils. Other fatty substances which are representative are the fatty alcohols (cetyl alcohol and stearyl alcohol), fatty acids (stearic acid) and waxes.

[0062] The emulsifiers are advantageously present in the compositions in a proportion ranging from 0.3 % to 30% by weight and preferably from 0.5 % to 30% by weight relative to the total weight of the composition.

[0063] In known fashion, the compositions of this invention may also contain adjuvants and additives that are common in the corresponding fields, such as hydrophilic or lipophilic gelling agents, preservatives, antioxidants, solvents, fragrances, fillers, UV-screening agents, dyestuffs, colorants, etc. Moreover, the subject compositions may contain hydrophilic or lipophilic bioaffecting active agents. The amounts of these various adjuvants, additives or active agents are those that are conventional in the cosmetic or pharmaceutical field, and, for example, range from 0.01 % to 20% of the total weight of the composition. Depending on their nature, these adjuvants, additives or active agents may be introduced into the fatty phase, into the aqueous phase and/or into lipid vesicles.

[0064] Among the active agents which the compositions of the invention may contain, particularly exemplary are the active agents which have an effect on the treatment of wrinkles or fine lines, other than manganese, and in particular keratolytic active agents. By the term “keratolytic” is intended an active agent which has desquamating, exfoliant or scrubbing properties, or an active agent capable of softening the horny layer.

[0065] Exemplary active agents for the treatment of wrinkles or fine lines, which the compositions of the invention may contain, include alverine or salts thereof, chlorine-channel openers, hydroxy acids and retinoids.

[0066] The hydroxy acids may be, for example, α -hydroxy acids or β -hydroxy acids, which may be linear, branched or cyclic, and saturated or unsaturated. The hydrogen atoms of the carbon chain may also be substituted with halogens, or halogenated alkyl, acyl, acyloxy, alkoxycarbonyl or alkoxy radicals having from 2 to 18 carbon atoms.

[0067] Exemplary hydroxy acids include, in particular, glycolic acid, lactic acid, maleic acid, tartaric acid, citric acid, 2-hydroxyalkanoic acid, mandelic acid, salicylic acid and alkyl derivatives thereof, for instance 5-n-octanoylsalicylic acid, 5-n-dodecanoylsalicylic acid, 5-n-decanoylsalicylic acid, 5-n-octylsalicylic acid, 5-n-heptyloxysalicylic acid or 4-n-heptyloxysalicylic acid and 2-hydroxy-3-methylbenzoic acid, or alkoxy derivatives thereof, for instance 2-hydroxy-3-methoxybenzoic acid.

[0068] And exemplary retinoids include, in particular, retinoic acid (all-trans or 13-cis) and derivatives thereof, retinol (vitamin A) and esters thereof, such as retinyl palmitate, retinyl acetate and retinyl propionate, and the salts thereof.

[0069] These active agents may be formulated, in particular, at concentrations ranging from 0.0001 % to 5 % by weight relative to the total weight of the composition.

[0070] The present invention also features a cosmetic regime/regimen for treating wrinkles and/or fine lines, comprising topically applying onto the skin a cosmetic composition containing an effective amount of manganese, formulated into a physiologically acceptable medium.

[0071] The cosmetic regime/regimen of the invention may be carried out, in particular, by topically applying the cosmetic composition as described above, via usual techniques. For example: application of creams, gels, sera, lotions,

makeup-removing milks or sunscreen compositions onto the skin or application of spray compositions.

[0072] In order to further illustrate the present invention and the advantages thereof, the following specific examples are given, it being understood that same are intended only as illustrative and in nowise limitative.

[0073] In said examples to follow, all parts and percentages are given by weight, unless otherwise indicated.

EXAMPLE 1:

[0074] Activity of manganese gluconate and of manganese chloride in a model of nerve/muscle junction (motor plate) obtained in a phrenic nerve/isolated diaphragm preparation: investigation of a myorelaxant effect:

[0075] The phrenic nerve and the diaphragm were carefully isolated and placed in a 50 ml tank filled with survival fluid (Krebs Henseleit liquid) maintained at a temperature of 37°C and oxygenated with a mixture of 95 % oxygen and 5% CO₂.

[0076] The variations in the tension of the diaphragm were then recorded with an initial preloading of several grams.

[0077] After a relaxation period of 30 min., the diaphragm was stimulated indirectly via the phrenic nerve.

[0078] On each preparation, the effect of the test products was evaluated, in a first stage, on the contractions induced by indirect stimulation via stimulation of the phrenic nerve (0.1 to 7 volts, 0.3 ms, 0.1 Hz) at increasing and cumulative concentrations from 10⁻⁶ M to 10⁻³ M.

[0079] The results obtained in the model of motor plate using the 2 manganese salts were as reported in the following Table:

TABLE:

Product	Concentration	% of inhibition (indirect stimulation)
Manganese gluconate (n=2)	10^{-3} M	25 %
MnCl ₂	10^{-3} M	10 %

EXAMPLES 2-6:

[0080] The following are specific examples of formulations according to the invention:

EXAMPLE 2:

[0081] Composition 1: Anti-wrinkle care lotion for the face:

Manganese gluconate	1.50 %
Antioxidant	0.05 %
Preservative	0.30 %
Ethanol (solvent)	8.00 %
Water	qs 100 %

[0082] This lotion acts on wrinkles during repeated use (application twice daily for one month).

EXAMPLE 3:

[0083] Composition 2: Care gel for the face:

Manganese chloride	0.50 %
Hydroxylpropylcellulose*	1.00 %
Preservative	0.30 %
Ethanol (solvent)	15.00 %
Antioxidant	0.05 %

Water	qs	100 %
-------	----	-------

*: Klucel H marketed by Hercules (gelling agent)

[0084] The gel obtained acts on wrinkles. It may be applied daily, morning and evening for one month.

EXAMPLE 4:

[0085] Composition 3: Care cream for the face (oil-in-water emulsion):

Manganese gluconate	0.50%
Glyceryl stearate (emulsifier)	2.00%
Polysorbate-60 (Tween 60 marketed by	
ICI) (emulsifier)	1.00%
Stearic acid	1.40%
Triethanolamine (neutralizer)	0.70%
Carbomer (Carbopol 940 marketed by	
Goodrich)	0.40%
Liquid fraction of karite butter	12.00%
Perhydrosqualene	12.00%
Preservative	0.30%
Fragrance	0.50%
Antioxidant	0.05%
Water	qs 100 %

[0086] A rich white cream is obtained, which reduces wrinkles and fine lines, and which may be applied daily.

EXAMPLE 5:

[0087] Composition 4: Care cream for the face (oil-in-water emulsion):

Manganese gluconate		0.10%
Glyceryl monostearate/distearate		2.00%
Cetyl alcohol		1.50%
Mixture of cetylstearyl alcohol/33 EO		
oxyethylenated cetylstearyl alcohol		7.00%
Polydimethylsiloxane		1.50%
Liquid petroleum jelly		17.50%
Preservative		0.30%
Fragrance		0.50%
Glycerol		12.50%
Water	qs	100 %

EXAMPLE 6:

[0088] Composition 5: Care cream for the face (oil-in-water emulsion):

Extract of walnut		5.00%
Glyceryl monostearate/distearate		2.00%
Cetyl alcohol		1.50%
Mixture of cetylstearyl alcohol/33 EO		
oxyethylenated cetylstearyl alcohol		7.00%
Polydimethylsiloxane		1.50%
Liquid petroleum jelly		17.50%
Preservative		0.30%
Fragrance		0.50%
Glycerol		12.50%

Water

qs

100%

[0089] While the invention has been described in terms of various specific and preferred embodiments, the skilled artisan will appreciate that various modifications, substitutions, omissions, and changes may be made without departing from the spirit thereof. Accordingly, it is intended that the scope of the present invention be limited solely by the scope of the following claims, including equivalents thereof.

0959334 051004
T081004